

GNFAC Avalanche Forecast for Wed Nov 28, 2018

Good Morning. This is Doug Chabot with early season avalanche and weather information from the Gallatin National Forest Avalanche Center on Wednesday, November 28th at 6:30 a.m. This bulletin is sponsored by [Spark R&D](#) and [World Boards](#). We will start issuing daily avalanche forecasts with [danger ratings](#) this Friday.

Mountain Weather

At 5 a.m. all the mountains except the Bridger Range are reporting snowfall. Big Sky to West Yellowstone has 3" with 1-2" in Hyalite and Cooke City. Mountain temperatures are near 20F and winds are 15-30 mph with gusts of 40 mph from the W-SW. Snowfall will add another 2-4" by tomorrow morning with Cooke City getting closer to 6". Winds will remain the same and temperatures will rise into the low 30s under mostly cloudy skies. Thursday looks to be partly cloudy with moderate SW winds and temperatures hitting the 30s.

Snowpack and Avalanche Discussion



Bridger Range Gallatin Range Madison Range Cooke City

Since Thanksgiving, over a foot of snow (1+” [SWE](#)) fell with strong west to southwest winds. Folks saw a few natural avalanches up Hyalite ([photo](#)), and in the Bridger Range ([report](#)), while skiers in Bear Basin in the northern Madison Range triggered a wind slab that they captured on [video](#). Avalanches were not widespread and were confined to the new snow. This matches our findings from the field which Alex describes in a [video from Middle Basin](#) (north of Big Sky) as well as [snowpit profiles](#) recently dug throughout our forecast area.

The underlying snowpack is generally strong and our primary concern is with today’s new snow. Snow and wind will create thick wind-drifts near the ridgelines which will be easily triggered. Recent avalanches or cracking ([photo](#)) of this new snow are signs of a rising avalanche danger. Stay out of avalanche terrain if you witness these warnings.



Lionhead Range

The snowpack in the Lionhead area outside West Yellowstone is shallow, weak and unstable. Alex was there on Monday and found a 2’ deep snowpack that consisted of sugary grains of facets capped by 6” of new snow ([photo](#)). His [video](#) shows the crumbly nature of the facets and their inability to support much new snow. With today’s snowfall and wind a skier or rider can trigger an avalanche, especially at the ridgelines where wind-drifts will be thick and unstable.

New this season, we added hyperlinks to the [Weather and Avalanche Log](#) and a new Menu item <[Avalanches and SnowPits](#)> with information on avalanche activity and incidents.

If you get out and have any avalanche or snowpack observations to share, contact us via our [website](#), email (mtavalanche@gmail.com), phone (406-587-6984), or Instagram ([#gnfacobs](#)).

Upcoming Avalanche Education and Events

Our education calendar is full of awareness lectures and field courses. Check it out: [Events and Education Calendar](#).

TONIGHT! November 29, 30 and Dec 1, 2 or 8, **Intro to Avalanche w/ Field Day**, [more info here](#)

December 1, [Klim Winter Kick-Off](#) and avalanche awareness, Rigby, Idaho

December 5, **1-hr Avalanche Awareness**, 6-7pm at REI, Bozeman

December 6, **1-hr Avalanche Awareness and Beacon Practice**, 6:30-8pm at Story Mill Park

COOKE CITY

Every Friday and Saturday, **Rescue Training and Snowpack Update**. Friday 6:30-7:30 p.m at the Soda Butte Lodge. Saturday anytime between 10-2 @ Round Lake.